39

40

41

42

43

44

45

46

47

48

49

50

51

52

53 54

55

49

50

51

52

53

54

55

56

KERALA MEDICAL JOURNAL

An Unusual case of Hemolytic Anemia

BL Harikrishnan, Ashwin K Hari, P Baburaj

Department of General Medicine, Jubilee Mission Medical College, Thrissur, Kerala*

ABSTRACT

Published on 27th September 2017

2

8

9

10

11

12

13

14 15

16 17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

Incidence of drug induced hemolysis is 12 to 18% of hemolytic anemia. Immune mediated and non immune mediated mechanisms are postulated for drug induced hemolytic anemia. We report a case of penicillin G induced hemolysis in a patient with RHD MS/ MR who was on penicillin prophylaxis, who presented with anemia and jaundice.

AQ1

*See End Note for complete author details

INTRODUCTION

Hemolytic anemia forms a major reason for recurrent jaundice with anemia. Manifestation depends on the etiology of the hemolysis. Broadly congenital and acquired causes are possible. Among the acquired auto-immune, both idiopathic and symptomatic like SLE, lymphoproliferative disease. Another important cause for acquired hemolytic anemia is drug induced; again both immune mediated and non immune mediated mechanisms are present. Here we report a case of Penicillin G induced immune mediated hemolytic anemia which is a very rare case.

CASE REPORT

A 35 years old female patient with RHD MS/MR was referred here for evaluation of recurrent jaundice. She is a known case of RHD MS/MR and was on Penicillin prophylaxis, oral Penicillin G. Jaundice started 2 years ago with recurrence. No itching, no discoloration of urine, no history of biliary colic.

Examination revealed icterus, mild pallor. No other general examination findings. Vitals normal, CVS examination revealed evidence of MS/MR, no CHF. No hepatosplenomegaly. Other systems were within normal limits.

Investigations- Total and differential leucocyte count: normal, ESR: 15mm/hr, Hb: 11gm%, Peripheral blood smear: normocytic normochromic RBC, no normoblasts or schiztocytes, reticulocyte count: 3%, LFT S. Bilirubin 4.5mg (4mg indirect and 0.5mg direct) SGOT and SGPT: normal, S.protein: normal, Coombs test: DCT positive ICT negative.

Considering immune hemolysis, she was investigated for the possibility of drug induced immune hemolysis. Since she was on regular oral Penicillin G, the test was directed on this basis. Penicillin adsorbed RBCs were treated with patients' serum which showed hemolysis. RBCs untreated with Penicillin G did not have hemolysis with patient's serum.

This confirmed the diagnosis of Penicillin induced Immune mediated hemolysis. On discontinuing penicillin prophylaxis features of hemolysis disappeared. Hb rose to 13gm%, reticulocyte count: 1.5%. Finally the diagnosis of Penicillin induced Immune mediated hemolysis was made

DISCUSSION

B lactum antibiotics Penicillin and Cephalosporins are known to produce immune mediated hemolysis. Drugs can produce hemolysis by immune mediated and non immune mediated mechanisms.

Immune mediated mechanisms

- · Revealing the masked antigen in RBCs, producing coombs positive hemolysis by a methyl dopa
- · Drug combining with RBC form new antigen thus antibody production and hemolysis - Penicillin type
- Immune complex, Innocent bystander type: Drug

Cite this article as: ---

Corresponding Author:

Kerala. Mobile: +919846569012 Email: hariclean@gmail.com

with antibody forming immune complex falls on RBC, producing hemolysis with complement activation.

Non immune mediated mechanisms

- 1. G6PD deficiency and other enzymopathy: hemolysis by drugs like Primaquine
- 2. Oxidation mechanism: Denatured Hb and RBC cell membrane producing Meth-hemoglobin and hemolysis; Drugs like Dapsone
- Non oxidative mechanisms: Proteins getting attached non immunologically to the RBC membrane; Drugs like Cephalothin, Cisplatin leading on to hemolysis.

END NOTE

Author Information

- Dr. BL Harikrishnan, Asst. Professor, Department of General Medicine, Jubilee Mission Medical College, Thrissur, Kerala
- 2. Dr. Ashwin K Hari, Senior Resident, Department of General Medicine, Jubilee Mission Medical

College, Thrissur, Kerala

3. Prof. Dr. P Baburaj, Department of General Medicine, Jubilee Mission Medical College, Thrissur, Kerala

Conflict of Interest: None declared

REFERENCES ???



- Kumar V, Abbas AK, Aster JC. Red blood cell and bleeding disorders. In: Kumar V, Abbas AK, Aster JC, eds. Robbins and Cotran Pathologic Basis of Disease. 9th ed. Philadelphia, PA: Elsevier Saunders; 2015: chap 14
- Michel M. Autoimmune and intravascular hemolytic anemias. In: Goldman L, Schafer AI, eds. Goldman-Cecil Medicine. 25th ed. Philadelphia, PA: Elsevier Saunders; 2016: chap 160
- 3. Price EA, Schrier SS. Extrinsic Nonimmune hemolytic anemias. In: Hoffman R, Benz EJ Jr, Silberstein LE, Heslop HE, Weitz JI, Anastasi, J, eds. *Hematology: Basic Principles and Practice*. 6th ed. Philadelphia, PA: Elsevier Saunders; 2013: chap 45.
- Petz LD, Garratty G. Immune Hemolytic Anemias. 2nd ed. Philadelphia: Churchill Livingstone; 2004:261–317.
- Johnson ST, Fueger JT, Gottschall JL. One center's experience: the serology and drugs associated with drug-induced immune hemolytic anemia – a new paradigm. Transfusion. 2007; 47: 697–702.
- Worlledge SM. Immune drug-induced hemolytic anemias. Semin Hematol. 1973; 10:327–344.

Author Queries???

AQ1: Kindly provide keywords AQ2: Please indicate references in text